



Short communication

'*Candidatus Rickettsia nicoyana*': A novel *Rickettsia* species isolated from *Ornithodoros kноxjonesi* in Costa Rica



Rolando D. Moreira-Soto^{a,b}, Andrés Moreira-Soto^a, Eugenia Corrales-Aguilar^a, Ólger Calderón-Arguedas^a, Adriana Troyo^{a,*}

^a Centro de Investigación en Enfermedades Tropicales (CIET), Facultad de Microbiología, Universidad de Costa Rica, San José, Costa Rica

^b Centro de Investigación en Estructuras Microscópicas (CIEMIC), Universidad de Costa Rica, San José, Costa Rica

ARTICLE INFO

Article history:

Received 11 May 2016

Received in revised form

14 December 2016

Accepted 26 February 2017

Available online 27 February 2017

Keywords:

Rickettsiales

Argasidae

Emballonuridae

Balantiopteryx

Costa Rica

ABSTRACT

Rickettsiae are intracellular bacteria commonly associated with hematophagous arthropods. Most of them have been described in hard ticks, but some have been found in soft ticks. Here we report the detection and isolation of a new *Rickettsia* from *Ornithodoros kноxjonesi* larvae collected from *Balantiopteryx plicata* (Emballonuridae) in Nicoya, Costa Rica. Two ticks were processed to detect *Rickettsia* spp. genes *gltA*, *ompA*, *ompB*, and *htrA* by PCR. Part of the macerate was also inoculated into Vero E6 and C6/36 cell lines, and cells were evaluated by Giménez stain, indirect immunofluorescence assay (IFA), and PCR. Both ticks were positive by PCR and rickettsial growth was successful in Vero E6 cells. Amplification and sequencing of near full length *rrs*, *gltA*, *sca4* genes, and fragments of *ompA* and *ompB* showed that the *Rickettsia* sp. was different from described species. The highest homologies were with '*Candidatus Rickettsia wissemanni*' and *Rickettsia peacockii*: 99.70% (1321/1325) with both sequences for *rrs*, 99.58% (1172/1177) and 99.76% (1246/1249) for *gltA*, 99.26% with both sequences (2948/2970 and 2957/2979) for *sca4*, 98.78% (485/491) and 98.39% (2069/2115) for *ompA*, and 98.58 (1453/1474) and 98.92% (1459/1475) for *ompB*; respectively. Bat blood, spleen, liver, and lung samples analyzed for *Rickettsia* detection were negative. Results demonstrate that the *Rickettsia* isolated from *O. kноxjonesi* is probably an undescribed species that belongs to the spotted fever group, for which '*Candidatus Rickettsia nicoyana*' is proposed. Considering that *B. plicata* inhabits areas where contact with humans may occur and that human parasitism by *Ornithodoros* has been reported in the country, it will be important to continue with the characterization of this species and its pathogenic potential.

© 2017 Elsevier GmbH. All rights reserved.

1. Introduction

The genus *Rickettsia* is comprised of obligate Gram-negative intracellular bacteria of arthropods (e.g., ticks, mites, fleas, and lice), some of which are considered endosymbionts while others are known pathogens of humans and other vertebrates (Rounds et al., 2012; Parola et al., 2013). In this respect, several rickettsiae that were initially considered non-pathogenic have been implicated in human infections and disease, and new species of *Rickettsia* continue to be identified in which their pathogenicity is yet to be determined (Parola et al., 2013).

Although the majority of known *Rickettsia* species have been described from hard ticks (Ixodida: Ixodidae), several species have

been found in association with soft ticks (Ixodida: Argasidae). For instance, *Rickettsia hoogstraalii* and similar genotypes have been detected in *Carios capensis*, and other species of *Carios*, *Argas*, and *Ornithodoros* in USA, Japan, Croatia, and several African countries (Cutler et al., 2006; Duh et al., 2010; Pader et al., 2012; Dietrich et al., 2014; Lafri et al., 2015). This species has also been detected in hard ticks (*Haemaphysalis* spp.) from Spain (Portillo et al., 2008). Additionally, *Rickettsia lusitaniae* was isolated from *Ornithodoros erraticus* from Portugal (Milhano et al., 2014), and '*Candidatus Rickettsia wissemanni*' was described from *Ornithodoros hasei* in French Guiana (Tahir et al., 2016). Other undescribed rickettsiae have also been found in *Argas* and *Carios* (Socolovschi et al., 2012; Loftis et al., 2005). The pathogenic potential of these recently described rickettsiae is unknown.

To date, there are no reports of rickettsiae associated with soft ticks from Central America. In Costa Rica, one study by Vargas in 1984 reports the occurrence of *Ornithodoros* associated with bats and refers to their importance since several people were bitten by

* Corresponding author at: Departamento de Parasitología, Facultad de Microbiología, Universidad de Costa Rica, San José, 11501, Costa Rica.

E-mail address: adriana.troyo@ucr.ac.cr (A. Troyo).

these ticks. However no analysis for *Rickettsia* presence was made (Vargas, 1984). There are more recent reports concerning distribution and human or animal parasitism by soft ticks in Central America, although none of them included the detection of rickettsiae (Rangel and Bermúdez-Castillero, 2013; Bermúdez et al., 2013; Bermúdez et al., 2015; Venzal et al., 2015). Therefore, the aim of this paper is to report the detection and isolation of a putative new species of *Rickettsia* from *Ornithodoros knoxjonesi* collected from *Balantiopteryx plicata* (Emballonuridae) bats in Costa Rica.

2. Materials and methods

2.1. Tick identification and detection of rickettsiae

Three soft tick larvae were collected in June 2014, from two *Balantiopteryx plicata* bats in Nicoya (10° 10' 10.342" N, 85° 26' 56.538" W), Guanacaste, Costa Rica. Bats were captured using mist nets surrounding their roosting habitat and collected following the recommendations of the Institutional Committee of Care and Use of Animals of the University of Costa Rica (CICUA-36-13), according to international animal welfare standards. For morphological evaluation of the larvae, one tick was cleared in lactophenol and mounted on a glass slide using Hoyer's medium according to established processing methods for mites (Krantz and Krantz, 2009); identification was made following keys for immature argasid ticks of the Neotropical Region (Barros-Battesti et al., 2013).

The two remaining ticks were processed individually to determine the presence of *Rickettsia* sp. First, the larvae were washed in 0.02% iodine and 70% ethanol solution for 10 min, then washed three times in sterile distilled water, macerated, and resuspended in 1 ml of brain-heart infusion broth (Hun et al., 2011). Genomic DNA was extracted from 200 µl of each macerate using Nucleospin Tissue[®] kit (Macherey-Nagel), following manufacturer's instructions. To detect *Rickettsia* spp. DNA, PCRs targeting four different genes were performed: *gltA* using primers CS-78 and CS-323 (401 bp product) (Labruna et al., 2004); *ompA* by a semi-nested PCR using primer pairs Rr190-70/Rr190-701 and Rr190-70/Rr190-602 (532 bp product) (Regnery et al., 1991; Roux et al., 1996); *ompB* using primers 120-M59f and 120-807 (856 bp product) (Roux and Raoult, 2000); and *htrA* by a nested PCR using primer pairs R17-122/R17-500 and TZ15/TZ16 (246 bp product) (Tzianabos et al., 1989; Blair et al., 2004). Blood, spleen, liver, and lungs from both bats were processed to detect the *Rickettsia gltA* and *ompB* gene fragments as stated previously. All the products were visualized on a 2% agarose gel after Gel Red staining, under UV light with a ChemiDocTM XRS+ System (BIO-RAD).

For molecular confirmation of tick species, extracted DNA was used to amplify the 16S mitochondrial rDNA gene using primers 16S+1 and 16S-1 (460 bp product), according to methods described previously (Black and Piesman, 1994; Mangold et al., 1998).

PCR products of ticks and rickettsiae were purified and sent to Macrogen Inc. (Seoul, South Korea) for sequencing with the corresponding primers (the second pair of primers was used in nested or semi-nested PCRs). DNA sequences were edited and assembled with DNA Baser (Heracle BioSoft S.R.L.), and a sequence homology search was performed against the NCBI database using BLAST.

2.2. Isolation of *Rickettsia* sp. in cell culture

Isolation was attempted from the macerate of the two tick larvae by inoculating 200 µl of the macerate into confluent monolayers of C6/36 (ATCC CRL-1660) and Vero E6 (ATCC CRL-1586) cell lines, according to previously described methods (Hun et al., 2011). Cells were maintained in RPMI or Minimal Essential Medium (MEM), respectively, supplemented with 5% fetal bovine serum, and

incubated at 28 °C in a 5% CO₂ atmosphere. The presence of *Rickettsia* spp. was monitored for 4 weeks by observing the appearance of a cytopathic effect and by Gimenez staining (Giménez, 1964). To confirm the identification of the isolates, DNA was extracted from positive cell cultures as stated above and PCR reactions were carried out to amplify the same four *Rickettsia* genes. The products were processed in the same manner and sequences were edited and compared to the ones obtained directly from the ticks.

Aliquots of the infected Vero E6 cells were treated according to methods previously described in order to prepare antigens on glass slides for testing by an indirect immunofluorescence assay (IFA) (Walker et al., 1977). Guinea pig sera reactive against *Rickettsia rickettsii* (end titer: 4 096) and *Rickettsia felis* (end titer: 512) were used at 2-fold serial dilutions to determine their end titers against the new *Rickettsia* isolate (Rivas et al., 2015). Anti-guinea pig IgG produced in goat and conjugated to fluorescein isothiocyanate (FITC) (Sigma–Aldrich) was used as secondary antibody at a dilution of 1:128. Negative antigen controls were included using Vero E6 mock infected cells.

2.3. Identification of *Rickettsia* sp.

To identify the species of *Rickettsia*, the *rrs*, *gltA*, and *sca4* genes were amplified from isolate DNA using primers previously described: 27F and 1492R for *rrs* (~1 465 bp sequence); CS2d, CSEndr, CS890r, and Rp877p for *gltA* (~1 270 bp sequence), and D1f, D928r, D767f, D1390r, D1219f, D1876r, D1738f, D2482r, D2338f, and D3069r for *sca4* (~3 065 bp sequence) (Lane, 1991; Roux et al., 1997; Mediannikov et al., 2004; Sekeyova et al., 2001). Additional *ompA* and *ompB* gene fragments were amplified with primers 190-70, 190-602, 190-3588, 190-4406, 190-4338, and 190-5238 for *ompA* (~532 and ~1 650 bp sequences); and 120-M59f, 120-807, 120-607, 120-1497, 120-1378, and 120-2399 for *ompB* (~1 565 bp sequence) (Regnery et al., 1991; Fournier et al., 1998; Roux and Raoult, 2000). PCR products were purified, sequenced, and assembled as stated above. For each gene, a sequence homology search was performed against the NCBI database using BLAST.

For phylogenetic analyses, sequences obtained in this study and those of described species of *Rickettsia* were aligned with GUIDANCE (Penn et al., 2010), using MAFFT algorithm, and a bootstrap of 100 repeats. From the resulting alignments, no columns or sequences were removed, and both ends of the alignments were trimmed with MEGA 5 (Tamura et al., 2011). From these trimmed alignments, the models of nucleotide substitutions were selected using jModelTest (Darrriba et al., 2012; Guindon and Gascuel, 2003). The GTR+G model was used for the *gltA*, *ompB*, *sca4* genes, and both trees of the *ompA* gene, the HKY+I model for the *rrs* gene, and the GTR+G model for a concatenated tree using *gltA*, *ompB*, *rrs*, and *sca4* gene fragments. The phylogenetic trees were constructed using a Bayesian MCMC phylogenetic reconstruction with Mr Bayes 3.2 software (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003). For constructing the trees, 4 chains were run, with a temperature of 0.2, for a maximum of 2,000,000 generations, discarding the initial 25%, and the standard deviation (SD) of split frequencies at the end of the runs were below 0.01 in all cases. Final editions of the trees were made using MEGA 5.

3. Results

Morphological characteristics placed tick larvae in the genus *Ornithodoros*. The partial sequence of the 16S mitochondrial rDNA gene confirmed the species as *Ornithodoros knoxjonesi* (KU663027).

Both ticks analyzed by PCR to detect *Rickettsia* spp. tested positive for the four genes (*gltA*, *ompA*, *ompB*, *htrA*). For all genes, sequences were identical between the two ticks, confirming that

Table 1
Highest BLAST sequence similarities of *rrs*, *gltA*, *sca4*, *ompA*, and *ompB* genes of the *Rickettsia* sp. from *O. knoxjonesi* against rickettsiae species in the NCBI database.

Gene fragment	% Sequence similarity (bp)	<i>Rickettsia</i> sp.
<i>rrs</i>	99.70 (1321/1325)	<i>Candidatus R. wissemanii</i> G1329 (LT558851.2)
	99.70 (1321/1325)	<i>R. peacockii</i> str. Rustic (CP001227.1)
	99.70 (1321/1325)	<i>R. rickettsii</i> str. Morgan (CP006010.1)
	99.62 (1320/1325)	<i>R. philipii</i> str. 364D (CP003308.1)
	99.62 (1320/1325)	<i>R. slovacica</i> str. D-CWPP (CP003375.1)
<i>gltA</i>	99.76 (1246/1249)	<i>R. peacockii</i> str. Rustic (CP001227.1)
	99.60 (1244/1249)	<i>R. philipii</i> str. 364D (CP003308.1)
	99.60 (1244/1249)	<i>R. slovacica</i> str. D-CWPP (CP003375.1)
	99.60 (1244/1249)	<i>R. parkeri</i> str. Portsmouth (CP003341.1)
	99.58 (1172/1177)	<i>Candidatus R. wissemanii</i> G1329 (LT558852.1)
	99.52 (1243/1249)	<i>R. rickettsii</i> str. Morgan (CP006010.1)
<i>sca4</i>	99.26 (2957/2979)	<i>R. peacockii</i> str. Rustic (CP001227.1)
	99.26 (2948/2970)	<i>Candidatus R. wissemanii</i> G1329 (LT558855.1)
	99.16 (2954/2979)	<i>R. rickettsii</i> str. Hlp#2 (CP003311.1)
	99.03 (2950/2979)	<i>R. philipii</i> str. 364D (CP003308.1)
	98.93 (2962/2994)	<i>R. slovacica</i> str. D-CWPP (CP003375.1)
<i>ompA</i> -Total*	98.78 (485/491)	<i>R. conorii</i> str. Malish 7 (AE006914.1)
	98.39 (2069/2115)	<i>Candidatus R. wissemanii</i> G1329 (LT558853.1)
	98.11 (2079/2113)	<i>R. philipii</i> str. 364D (CP003308.1)
	98.30 (2073/2113)	<i>R. rickettsii</i> str. R (CP006009.1)
	97.83 (2077/2113)	<i>R. slovacica</i> str. D-CWPP (CP003375.1)
<i>ompB</i>	98.92 (1459/1475)	<i>R. peacockii</i> str. Rustic (CP001227.1)
	98.58 (1453/1474)	<i>Candidatus R. wissemanii</i> G1329 (LT558854.1)
	98.24 (1449/1475)	<i>R. philipii</i> str. 364D (CP003308.1)
	98.24 (1449/1475)	<i>R. rickettsii</i> str. Hlp#2 (CP003311.1)
	97.83 (1443/1475)	<i>R. slovacica</i> str. D-CWPP (CP003375.1)

*Values represent a sum of sequence similarity of both gene fragments.

they harbored the same species of *Rickettsia*. Blood, spleen, liver, and lung samples from both bats processed showed no presence of *Rickettsia* DNA.

The *Rickettsia* sp. was successfully cultured in Vero E6 cells from both ticks, and then confirmed through specific PCR and IFA. The reaction observed in the IFA was stronger with guinea pig serum reactive to *R. rickettsii* (Fig. S1 in Supplementary materials); end titer to the *Rickettsia* sp. isolate was similar to the end titer obtained with *R. rickettsii* antigen (en titer = 4 096) and higher than the end titer to *R. felis* (en titer < 128). Guinea pig serum reactive to *R. felis* (end titer = 512) had an end titer of 256 to the *Rickettsia* sp. isolate and of 128 to *R. rickettsii* antigen. All partial gene sequences obtained from the two isolates were identical to the ones obtained directly from the ticks. Preliminary analysis of these partial sequences determined that they were most similar to sequences of rickettsiae from the spotted fever group (SFG), but presented differences with those of described species.

Almost complete sequences of *rrs* (1 325 bp), *gltA* (1 249 bp), and *sca4* (2 979 bp) were obtained, as well as additional fragments of *ompA* (2 115 bp total) and *ompB* (1 475 bp). As for described species, sequences showed higher similarity with *Rickettsia peacockii* (Table 1): 99.70% (1321/1325) for *rrs*, 99.76% (1246/1249) for *gltA*, 99.26% (2957/2979) for *sca4*, 97.83% (2077/2113) for *ompA*, and 98.92% (1459/1475) for *ompB*. However, some sequences were most similar to the recently reported '*Candidatus Rickettsia wissemanii*' (*Rickettsia* sp. G1329): 99.70% (1321/1325) for *rrs*, 99.58% (1172/1177) for *gltA* (bp), 99.26% (2948/2970) for *sca4* (bp), and 98.58% (1453/1474) for *ompB* (Table 1). The *ompA* sequence from *Rickettsia* sp. in this study and '*Candidatus Rickettsia wissemanii*' were comparable along a shorter fragment of 491 bp, which showed 98.78% (458/491) similarity.

Phylogenetic analyses of *rrs*, *gltA*, *sca4*, *ompA*, and *ompB* confirmed that the *Rickettsia* isolated from *O. knoxjonesi* groups with *Rickettsia peacockii* and '*Candidatus R. wissemanii*', within the SFG (Figs. S2 and S3 in Supplementary materials). Fig. 1 shows the

concatenated tree using *rrs*, *gltA*, *sca4*, and *ompB* sequences (*ompA* was excluded because the sequence of '*Candidatus R. wissemanii*' was much shorter than all others).

These results suggest that the *Rickettsia* obtained from *O. knoxjonesi* probably corresponds to a new species (Raoult et al., 2005), and the name '*Candidatus Rickettsia nicoyana*' is proposed until further characterization of the isolates is performed.

DNA sequences of '*Candidatus R. nicoyana*' are deposited in GenBank under accession numbers KX228147 (*rrs*), KX228143 (*gltA*), KY296503 (*sca4*), KX228144 and KY296502 (*ompA*), KX228145 (*ompB*), and KX228146 (*htrA*). Isolates are stored in the Virology Laboratory of the Research Center for Tropical Diseases (Centro de Investigación en Enfermedades Tropicales, CIET) Faculty of Microbiology, University of Costa Rica.

4. Discussion and conclusions

The sequence analyses of the four genes revealed that this putative new species, provisionally named "*Candidatus R. nicoyana*", belongs to the spotted fever group (SFG), specifically to the subgroup that contains *Rickettsia rickettsii*. Since all of the analyses confirm that it does not group with *R. felis*, it is evidently different from most of the species known to be associated with Argasidae, such as *R. hoogstraalii* and *Rickettsia lusitaniae* (Milhano et al., 2014; Duh et al., 2010).

The sequence comparisons using *rrs*, *gltA*, *sca4*, *ompA*, and *ompB* genes suggest that the putative new *Rickettsia* found in this study is closely related to '*Candidatus R. wissemanii*', which is a newly described rickettsia detected in the bat tick *O. hasei* (Tahir et al., 2016). Although sequence similarity between '*Candidatus R. nicoyana*' and '*Candidatus R. wissemanii*' in the *sca4* gene was very close to the cutoff value used to separate species of *Rickettsia* (99.26%), differences in *rrs*, *gltA*, and *ompB* sequences separate '*Candidatus R. nicoyana*' from known species (Fournier et al., 2003; Raoult et al., 2005). Unfortunately, the partial *ompA* sequence available for




Fig. 1. Phylogenetic analysis based on the concatenated *gltA*, *ompB*, *rrs*, and *sca4* gene fragments of *Rickettsia* spp. The numbers on the branches indicate posterior probabilities.

'*Candidatus R. wissemanni*' is much shorter than the one of '*Candidatus R. nicoyana*', so further conclusions are not possible for this gene. Furthermore, a spotted fever group *Rickettsia* sp. that was detected in *Carios kelleyi* (= *Ornithodoros kelleyi*) was not characterized further and there are no corresponding sequences for *rrs*, *sca4*, *gltA*, or *ompB* for comparisons (Loftis et al., 2005). Overall, it is not surprising to find a close phylogenetic relationship between rickettsiae in *Ornithodoros* ticks in the Neotropical Region, with genetic divergence as a consequence of tick speciation in different geographical areas.

Although no rickettsiae were detected in blood or tissues of the bats analyzed in this study, there is evidence that supports infection in bats. In a recent study, Dietrich et al. (2016) detected the *gltA* fragment of a *Rickettsia* sp. closely related to *Rickettsia conorii* in bat blood samples. Moreover, seropositivity to SFG rickettsiae has been reported in several bat species from Brazil, most being reactive to two or more SFG *Rickettsia* antigens (D'Auria et al., 2010), which may be the result of cross reactivity between SFG rickettsiae, as was observed in the present study. More investigations are needed to determine potential host-pathogen relationships between these rickettsiae and bats, and the possible role of bats in their transmission cycle.

The ticks *O. knoxjonesi* were collected in a *B. plicata*, and this bat species is known to roost in colonies where human contact is possible (Nagy et al., 2014). Moreover, '*Candidatus R. nicoyana*' was able to infect Vero cells, which is a cell line derived from primate kidney epithelium that can sustain the growth of pathogenic rickettsiae of the same subgroup such as *R. rickettsii* and *R. philipii* (Philip et al., 1981; Hun et al., 2008). Considering that human parasitism with *Ornithodoros* sp. has been previously reported in Costa Rica (Vargas, 1984), the importance of characterizing this putative new species and its pathogenic potential is evident. In this sense, it has been said that argasid ticks could play a role in the transmission of pathogenic *Rickettsia* to humans and that bats may serve as reservoirs for bacterial zoonoses (Loftis et al., 2005). Further studies are needed to determine the distribution, natural transmission cycle, and pathogenic potential to both bats and humans of this *Rickettsia* isolated.

Acknowledgements

Authors thank José Venzal for help with tick identification, Francisco Vega for technical assistance in IFAs and isolation of rickettsiae, Amanda Vicente for collaborating in bat captures and blood collection, Andrea Rodriguez, Kristel Murillo, and Dione Palma for helping with DNA extractions and PCRs, and Djamel Tahir for making sequences of '*Candidatus R. wissemanni*' (*Rickettsia* sp.

G1329) available for comparison. Financial support was provided by University of Costa Rica projects 803-B4-055 and FEES-CONARE 803-B4-656.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ttbdis.2017.02.015>.

References

- Barros-Battesti, D.M., Ramirez, D.G., Landulfo, G.A., Faccini, J.L., Dantas-Torres, F., Labruna, M.B., Venzal, J.M., Onofrio, V.C., 2013. Immature argasid ticks: diagnosis and keys for Neotropical region. *Rev. Bras. Parasitol. Vet.* 22, 443–456.
- Bermúdez, S.C.E., Miranda, R.J., Kadoch, N., 2013. Reporte de larvas de *Ornithodoros puertoricensis* Fox 1947 (Ixodida: Argasidae) parasitando a *Rhinella marina* (L. 1758) (Anura: Bufonidae) en David Chiriquí, Panamá. *Puente Biológico* 5, 81–85.
- Bermúdez, S., Miranda, R.J., Cleghorn, J., Venzal, J.M., 2015. *Ornithodoros (Alectorobius) puertoricensis* (Ixodida: Argasidae) parasitizing exotic reptiles pets in Panamá. *FAVE Secc. Cienc. Vet.* 14, 1–5.
- Black, W.C., Piesman, J., 1994. Phylogeny of hard- and soft-tick taxa (Acari: ixodida) based on mitochondrial 16S rDNA sequences. *Proc. Natl. Acad. Sci. U. S. A.* 91, 10034–10038.
- Blair, P.J., Jiang, J., Schoeler, G.B., Moron, C., Anaya, E., Cespedes, M., Cruz, C., Felices, V., Guevara, C., Mendoza, L., Villaseca, P., Sumner, J.W., Richards, A.L., Olson, J.G., 2004. Characterization of spotted fever group *Rickettsia* in flea and tick specimens from Northern Peru. *J. Clin. Microbiol.* 42, 4961–4967.
- Cutler, S.J., Browning, P., Scott, J.C., 2006. *Ornithodoros moubat*, a soft tick vector for *Rickettsia* in East Africa? *Ann. N. Y. Acad. Sci.* 44, 373–377.
- D'Auria, S.R.N., Camargo, M.C.G.O., Pacheco, R.C., Savani, E.S.M.M., Dias, M.A.G., Da Rosa, A.R., De Almeida, M.F., Labruna, M.B., 2010. Serologic survey for rickettsiosis in bats from São Paulo city, Brazil. *Vector Borne Zoonotic Dis.* 10, 459–463.
- Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods* 9, 772.
- Dietrich, M., Lebarbenchon, C., Jaeger, A., Rouzic, C.L., Bastien, M., Lagadec, E., McCoy, K.D., Pascalis, H., Corre, M.L., Dellagi, K., Tortosa, P., 2014. *Rickettsia* spp. in seabird ticks from western Indian ocean islands, 2011–2012. *Emerg. Infect. Dis.* 20, 838–842.
- Dietrich, M., Tjale, M.A., Weyer, J., Kearney, T., Seemark, E.C.J., Nel, L.H., Monadjem, A., Markotter, W., 2016. Diversity of *Bartonella* and *Rickettsia* spp. in bats and their blood-feeding ectoparasites from South Africa and Swaziland. *PLoS One* 11, e0152077.
- Duh, D., Punda-Polic, V., Avsic-Zupanc, T., Bouyer, D., Walker, D.H., Popov, V.L., Jelovsek, M., Gracner, M., Trilar, T., Bradaric, N., Kurtti, T.J., Strus, J., 2010. *Rickettsia hoogstraalii* sp. nov.: isolated from hard- and soft-bodied ticks. *Int. J. Syst. Evol. Microbiol.* 60, 977–984.
- Fournier, P.E., Roux, V., Raoult, D., 1998. Phylogenetic analysis of spotted fever group rickettsiae by study of the outer surface protein rOmpA. *Int. J. Syst. Bacteriol.* 48, 839–849.
- Fournier, P.E., Dumler, J.S., Greub, G., Zhang, J., Wu, Y., Raoult, D., 2003. Gene sequence-based criteria for identification of new *Rickettsia* isolates and description of *Rickettsia heilongjiangensis* sp. nov. *J. Clin. Microbiol.* 41, 5456–5465.
- Giménez, D.F., 1964. Staining rickettsiae in yolk-sac cultures. *Stain Technol.* 39, 135–140.

- Guindon, S., Gascuel, O., 2003. A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Syst. Biol.* 52, 696–704.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 17, 754–755.
- Hun, L., Cortés, X., Taylor, L., 2008. Molecular characterization of *Rickettsia rickettsii* isolated from human clinical samples and from the rabbit tick *Haemaphysalis leporispalustris* collected at different geographic zones in Costa Rica. *Am. J. Trop. Med. Hyg.* 79, 899–902.
- Hun, L., Troyo, A., Taylor, L., Barbieri, A.M., Labruna, M.B., 2011. First report of the isolation and molecular characterization of *Rickettsia amblyommii* and *Rickettsia felis* in Central America. *Vector Borne Zoonotic Dis.* 11, 1395–1397.
- Krantz, D.E., Krantz, G.W., 2009. Collecting, rearing, and preparing specimens. In: Krantz, G.W., Walter, D.E. (Eds.), *A Manual of Acarology*. Tech University Press, Texas, pp. 83–96.
- Labruna, M.B., Whitworth, T., Horta, M.C., Bouyer, D.H., MacBride, J.W., Pinter, A., Popov, V., Gennari, S.M., Walker, D.H., 2004. *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the State of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J. Clin. Microbiol.* 42, 90–98.
- Lafri, I., Leulmi, H., Baziz-Neffah, F., Lalout, R., Mohamed, C., Mohamed, K., Parola, P., Bitam, I., 2015. Detection of a novel *Rickettsia* sp. in soft ticks (Acari Argasidae) in Algeria. *Microbes Infect.* 17, 859–861.
- Lane, D.J., 1991. 16S/23S rRNA sequencing, p 115–147. In: Stackebrandt, E., Goodfellow, M. (Eds.), *Nucleic Acids Techniques in Bacterial Systematics*. John Wiley & Sons, Chichester.
- Loftis, A.D., Gill, J.S., Schrieffer, M.E., Levin, M.L., Eremeeva, M.E., Gilchrist, M.J.R., Dasch, G.A., 2005. Detection of *Rickettsia*, *Borrelia*, and *Bartonella* in *Carios kelleyi* (Acari: Argasidae). *J. Med. Entomol.* 42, 473–480.
- Mangold, A.J., Bargaes, M.D., Mas-Coma, S., 1998. Mitochondrial 16S rDNA sequences and phylogenetic relationships of species of *Rhipicephalus* and other tick genera among Metastriata (Acari: Ixodidae). *Parasitol. Res.* 84, 478–484.
- Mediannikov, O.Y., Sidelnikov, Y., Ivanov, L., Mokretsova, E., Fournier, P.E., Tarasevich, I., Raoult, D., 2004. Acute tick-borne rickettsiosis caused by *Rickettsia heilongjiangensis* in Russian Far East. *Emerg. Infect. Dis.* 10, 810–817.
- Milhano, N., Palma, M., Marcili, A., Nuncio, M.S., de Carvalho, I.L., De Sousa, R., 2014. *Rickettsia lusitaniae* sp. nov. isolated from the soft tick *Ornithodoros erraticus* (Acarina: Argasidae). *Comp. Immunol. Microbiol. Infect. Dis.* 37, 189–193.
- Nagy, M., Knörnschild, M., Günther, L., Mayer, F., 2014. Dispersal and social organization in the Neotropical Grey sac-winged bat *Balantiopteryx plicata*. *Behav. Ecol. Sociobiol.* 68, 891–901.
- Pader, V., Buniak, J.N., Abdissa, A., Adamu, H., Tolosa, T., Gashaw, A., Cutler, R.R., Cutler, S.J., 2012. *Candidatus Rickettsia hoogstraalii* in Ethiopian *Argas persicus* ticks. *Ticks Tick Borne Dis.* 3, 337–344.
- Parola, P., Paddock, C.D., Socolovschi, C., Labruna, M.B., Mediannikov, O., Kernif, T., Abdad, M.Y., Stenos, J., Bitam, I., Fournier, P., Raoult, D., 2013. Update on tick-borne rickettsioses around the world: a geographic approach. *Clin. Microbiol. Rev.* 26, 657–702.
- Penn, O., Privman, E., Ashkenazy, H., Landan, G., Graur, D., Pupko, T., 2010. GUIDANCE: a web server for assessing alignment confidence scores. *Nucleic Acids Res.* 38, W23–W28.
- Philip, R.N., Lane, R.S., Casper, E.A., 1981. Serotypes of tick-borne spotted fever group rickettsiae from western California. *Am. J. Trop. Med. Hyg.* 30, 722–727.
- Portillo, A., Santibanez, P., Santibanez, S., Perez-Martinez, L., Oteo, J.A., 2008. Detection of *Rickettsia* spp. in *Haemaphysalis* ticks collected in La Rioja, Spain. *Vector Borne Zoonotic Dis.* 8, 653–658.
- Rangel, G., Bermudez-Castillero, S.E., 2013. Nota sobre un caso de parasitismo de *Ornithodoros* sp. (Ixodida: Argasidae) en una mujer proveniente de La Laja Los Santos, Panamá. *Rev. Med. Panamá* 34, 37–39.
- Raoult, D., Fournier, P.E., Eremeeva, M., Kelly, P.J., Oteo, J.A., Sekeyova, Z., Tamura, A., Tarasevich, I., Zhang, L., 2005. Naming of rickettsiae and rickettsial diseases. *Ann. N. Y. Acad. Sci.* 1063, 1–12.
- Regnery, R.L., Spruill, C.L., Plikaytis, B.D., 1991. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J. Bacteriol.* 173, 1576–1589.
- Rivas, J.J., Moreira-Soto, A., Alvarado, G., Taylor, L., Calderón-Arguedas, O., Hun, L., Corrales-Aguilar, E., Morales, J.A., Troyo, A., 2015. Pathogenic potential of a Costa Rican strain of ‘*Candidatus Rickettsia amblyommii*’ in guinea pigs (*Cavia porcellus*) and protective immunity against *Rickettsia rickettsii*. *Ticks Tick-Borne Dis.* 6, 805–811.
- Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Rounds, M.A., Crowder, C.D., Matthews, H.E., Philipson, C.A., Scoles, G.A., Ecker, D.J., Schutzer, S.E., Eshoo, M.W., 2012. Identification of endosymbionts in ticks by broad-range polymerase chain reaction and electrospray ionization mass spectrometry. *J. Med. Entomol.* 49, 843–850.
- Roux, V., Raoult, D., 2000. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int. J. Syst. Evol. Microbiol.* 50, 1449–1455.
- Roux, V., Fournier, P.E., Raoult, D., 1996. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J. Clin. Microbiol.* 34, 2058–2065.
- Roux, V., Rydkina, E., Eremeeva, M., Raoult, D., 1997. Citrate synthase gene comparison, a new tool for phylogenetic analysis, and its application for the rickettsiae. *Int. J. Syst. Bacteriol.* 47, 252–261.
- Sekeyova, Z., Roux, V., Raoult, D., 2001. Phylogeny of *Rickettsia* spp. inferred by comparing sequences of ‘gene D’, which encodes an intracytoplasmic protein. *Int. J. Syst. Evol. Microbiol.* 51, 1353–1360.
- Socolovschi, C., Kernif, T., Raoult, D., Parola, P., 2012. *Borrelia*, *Rickettsia*, and *Ehrlichia* species in bat ticks, France, 2010. *Emerg. Infect. Dis.* 18, 1966–1975.
- Tahir, D., Socolovschi, C., Marié, J.L., Ganay, G., Berenger, J.M., Bompar, J.M., Blanchet, D., Cheuret, M., Mediannikov, O., Raoult, D., Davoust, B., Parola, P., 2016. New *Rickettsia* species in soft ticks *Ornithodoros hasei* collected from bats in French Guiana. *Ticks Tick Borne Dis.* 7, 1089–1096.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5 Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Tzianabos, T., Anderson, B.E., Mcdade, J.E., 1989. Detection of *Rickettsia rickettsii* DNA in clinical specimens by using polymerase chain reaction technology. *J. Clin. Microbiol.* 27, 2866–2868.
- Vargas, V.M., 1984. Occurrence of the bat tick *Ornithodoros (Alectorobius) kelleyi* Cooley & Kohls (Acari: Argasidae) in Costa Rica and its relation to human bites. *Rev. Biol. Trop.* 32, 103–107.
- Venzal, J.M., Hernández, L.V., Labruna, M.B., Nava, S., 2015. *Nothoaspis reddelli* Keirans & Clifford, 1975 (Acari, Ixodida, Argasidae): distribution extension. *Check List* 11, 1698.
- Walker, D.H., Harrison, A., Henderson, F., Murphy, F.A., 1977. Identification of *Rickettsia rickettsii* in a guinea pig model by immunofluorescent and electron microscopic techniques. *Am. J. Pathol.* 86, 343–358.